

A Case Report**Penile Thread Tourniquet Syndrome: A Case Report and Review of Age-Related Variability**Sadaqat Ullah Rehmat¹, Jawad khan², Attia Mahmood³, Asifa Irfan⁴, Aisha Habib⁵¹⁻⁵ Department of Urology Ayub Teaching Hospital Abbottabad**ABSTRACT**

Penile Thread Tourniquet Syndrome (PTTS), a rare variant of Penile Tourniquet Syndrome (PTS), is a paediatric urological emergency characterized by external constriction of penis by hair or thread, often leading to necrosis, ischemia, and oedema. A urethro-cutaneous fistula, urethral transection, and penile amputation are possible outcomes of this disorder if left untreated. Though commonly reported in infants, this case details a rare occurrence in a 10-year-old male. A circumferential thread at the penoscrotal junction caused localized infection and oedema without urethral damage. Surgical intervention and conservative management resulted in a favourable outcome. This case emphasizes the need for clinical vigilance in atypical age groups and rare causes of penile injury

Keywords: Penile thread torniquet syndrome (PTTS), Penile torniquet syndrome (PTS), Penile injury, Thread strangulation

INTRODUCTION

Penile tourniquet syndrome (PTS) is a very rare condition typically attributed to a hair coil wrapped around the sulcus coronarius of the penis. It occurs predominantly in circumcised boys aged 0-6 years [1-2]. Ischaemia necrosis, urethrocutaneous fistulas, gangrene, and even auto-amputation are among the most dangerous side effects of PTS[3]. Though hair tourniquets have also been observed on the clitoris, labia, and digits, this phenomenon primarily affects male circumcised individuals[4]. This report highlights a rare case of thread-induced PTS in a 10-year-old male, which is a less typical age for presentation.

Case presentation

A 10-year-old circumcised male presented to the Emergency Department at Ayub Teaching Hospital, Abbottabad, with penile swelling, redness, and mild pain. The patient's father claims that he experienced penile swelling, which the parents unintentionally noticed while changing his clothes because the child did not initially report any symptoms. The patient was unaware of what was causing his symptoms. On frequent inquiry, it was after more investigation, no history of trauma, insect bites, or comparable incidents was found. He reported normal urinary voiding. Initially assessed by a general practitioner (GP) in the local area, the swelling was misdiagnosed as an allergic reaction. Despite receiving prescribed medications, there was no improvement in symptoms. Therefore, the patient was referred to a Tertiary care hospital. Upon careful examination by the Urology team, a thread was found tightly wrapped around the penoscrotal junction, leading to oedematous swelling and a circumscribed laceration at the base of the penis. The area appeared inflamed, and purulent discharge was noted, indicating an ongoing infection. The patient was immediately shifted to the operating theatre, where the thread was carefully removed and dead tissue debrided under local anaesthetic. (Figure 1). Patient was catheterized and at the time, there was no urethral involvement appreciated. Wound care was initiated, including daily dressing with povidone-iodine and normal saline. Intravenous antibiotics and analgesics were administered to manage infection and pain. The patient was closely monitored for signs of tissue viability and possible complications such as necrosis, urethral involvement, or persistent swelling. The patient's wound showed significant improvement with conservative management. He was discharged home with instructions for continued wound care, oral antibiotics, and analgesics. A follow-up appointment was scheduled after three weeks to ensure complete healing and to assess for any long-term complications. The patient's father denied any involvement of an external culprit in causing the condition, and the child was assessed to be psychologically fit. Follow-up after 3 weeks, wound showed significant healing (Figure 2). No signs of infection or pus present; catheter removed, patient was able to pass urine spontaneously and sent to home.

How To Cite This Article: Rehmat SU, Khan J, Mahmood A, Irfan A, Habib A. Penile Thread Tourniquet Syndrome: A Case Report and Review of Age-Related Variability. *Pak J Urol.* 2025;3(2):48–50.

Corresponding Author: Jawad khan

Assistant Prof Department of Urology Ayub
Teaching Hospital Abbottabad

Email: jawadkhanjadoon77@gmail.com

ORCID: <https://orcid.org/0000-0002-0852-4840>

ARTICLE TRACKING

Received:	22- JULY -2025
Revision:	26-AUG-2025
Accepted:	23-SEP-2025
Published:	10-JAN- 2025
DOI:	10.69885/pju.v4i1.130

Figure 1: Post-Operative image of an infected circumferential wound



Figure 2: Follow up after 3 weeks of a wound healing



Discussion

PTS commonly affects an infant's genitalia, toes, and fingers, among other appendicular organs [5]. Over 90% of postpartum women experience telogen effluvium, a condition that causes significant hair loss [6]. The fact that our case reports a 10-year-old, highlights that it may be overlooked in older children. Increased independence in hygiene and dressing can delay detection by parents, contributing to late presentation. Several case reports have been published on this condition; however, very few have been reported from Pakistan [7]. Nonetheless, our case offers a number of unusual characteristics that can deepen our clinical knowledge of this condition. coronal sulcus as the common site for hair entrapment is often emphasised in the literature, very few, including our case, report a constriction at the base of the penile shaft, suggesting that the anatomical vulnerability of the coronal sulcus is not a prerequisite for the development of PTS [8]. Approximately 79% of documented cases use hair as the restricting substance. However, our case adds to the small pool of such recorded cases of thread-induced PTS, which have been reported seldom. Child abuse must be taken into account in certain

circumstances. According to reports, children with PTS may purposefully or compulsively wrap things around their genitalia in an attempt to regulate nocturnal enuresis, which may be linked to underlying psychological or behavioral issues [9]. Our patient had no known history of mental illness or behavioral disorders. The only relevant history was of nocturnal enuresis, but there was no evidence, either admitted or observed, of intentional thread wrapping by the child or family members. Simple cutting or unwinding of the restricting material is used to manage early cases. Surgery, however, can be necessary in cases of more advanced presentation with significant tissue damage or infection [10]. Furthermore, careful inspection in well-lit conditions and occasionally with magnification to identify the fibers is of paramount importance, as evident from this case.

Conclusion

PTTS is a urological emergency that, although rare, can lead to serious complications if not promptly identified and managed. Since prompt medical intervention for any penile anomaly can stop progression, given that situations like these might go unnoticed at the primary care level, general practitioners, especially those in outlying areas, should consider this diagnosis, especially when working with low socioeconomic communities. This case underlines the importance of prompt recognition, thorough examination, and appropriate follow-up in pediatric patients.

Consent:

Written informed consent was obtained from patient's parents who participated in this case.

Authors Contribution

Sadaqat Ullah Rehmat¹: Conceptualization, Methodology, Software

Jawad.khan²: Data curation, Writing- Original draft preparation.

Attia Mahmood³: Visualization, Investigation.

Asifa Irfan⁴ & **Aisha Habib**⁵: Writing- Reviewing and Editing.

References

1. Adjei NN, Lynn AY, Grimshaw A, Brault MA, Dugan K, Vash-Margita A. Systematic Literature Review of Pediatric Male and Female Genital Hair Thread Tourniquet Syndrome. *Pediatric emergency care*.2022;38:e799.e804. doi.org/10.1097/pec.0000000000002383.
2. AlJahdali EA. The First Report of Isolated Clitoral Hood Hair-Thread Tourniquet Syndrome: A Study of Six Patients. *Cureus*.2022;14:e26472. doi.org/10.7759/cureus.26472.
3. Djokic D, Milani GP, Lava SAG, Gualco G, Corigliano

T, Bianchetti MG, et al. Hair-thread strangulation syndrome in childhood: a systematic review. Swiss medical weekly. 2023;153:40124. doi.org/10.57187/smw.2023.40124.

4. Holmes A, Tissot S, Braslis K. Case report: Challenging adult case of erosive coronal sulcus hair thread tourniquet. Urology case reports. 2022;41:101983. doi.org/10.1016/j.eucr.2021.101983.

5. Ibrahim A, Abiir Y, Gounni Y, Kallat A, Boualaoui I, El Abidi H. Thread-induced penile injury in a child: Surgical management and outcomes in a hair-thread tourniquet syndrome. International journal of surgery case reports. 2025;135:111780. doi.org/10.1016/j.ijscr.2025.111780.

6. Koushik P, Daniel B, Meyyappan V, Sekar H, Krishnamoorthy S. A Case Series on Penile Garroting: Causes, Considerations, and Consequences. Cureus. 2024;16:e70190. doi.org/10.7759/cureus.70190.

7. Low P, Wang L, Li KD, Shibley WP, Cedars BE, Holler JT, et al. Thematic analysis of the psycho-sexual symptoms in patients with Peyronie's disease present on online forums. International journal of impotence research. 2023;35:533-541. doi.org/10.1038/s41443-022-00589-x.

8. Raheem OA, Kalidoss S, Pomar HA, Suarez- Sarmiento A, Jr., Perito P. Double distal corporal anchoring stitch for lateral penile implant cylinder extrusion. The journal of sexual medicine. 2024;21:96770. doi.org/10.1093/jsxmed/qdae100.

9. Rampersad B, Dubay C, Doodnath R. Penile hair tourniquet to base of penis resulting in circumferential separation of shaft skin without vascular compromise: An unusual case report. Urology case reports. 2022;42:102030. doi.org/10.1016/j.eucr.2022.102030.

10. Reichenbach A, Gayrel P, Morel Journal N, Carnicelli D, Vedrine N, Charbonnel C, et al. Surgical management for penile fracture: A retrospective, observational, multicenter study. The French journal of urology. 2024;34:102590. doi.org/10.1016/j.fjurol.2024.102590.



Licensing and Copyright Statement

All articles published in the Pakistan Journal of Urology are licensed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0). This license allows users to share (copy and redistribute) and adapt (remix, transform, and build upon) the published material for any purpose, including commercial, provided appropriate credit is given to the original author(s) and the source (Pakistan Journal of Urology), link to the license is provided, and any changes made are indicated. This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/). © The Author(s) 2025.

Publisher: Institute of Kidney Diseases and Pakistan Association of Urological Surgeons (PAUS)