PAKISTAN JOURNAL OF UROLOGY

ISSN: 3005-7582 (Online) : ISSN: 3005-7574 (Print)

OPEN ACCESS

Pakistan J Urol 2024, 02(1): P:114-117

A Case Report

Recurrent Para - Testicular Rhabdomyosarcoma A Case Report

Sana Hamayun¹, Wagma Hidayat², Hamza Rafi³, Haq Nawaz⁴ ^{1,2,3,4}Department of General Surgery, Hayatabad medical complex, Peshawar, Pakistan.

ABSTRACT

Background: Para-testicular RMS is not frequently reported and originates from the stromal tissues of the spermatic cord, epididymis, testis and tunica. It is a very uncommon feature in RMS and may be seen in a small percentage of patients and most often in children. At times the diagnosis may pose as a problem and physical examination may be inadequate at which time ultrasonography comes in handy. The main form of management is surgical through radical orchiectomy and the medical form is through Actinomycin D, vincristine and cyclophosphamide and very rarely radiotherapy.

Keywords: Para-testicular rhabdomyosarcoma, Radical orchiectomy, Chemotherapy, Ultrasonography

Introduction:

There are following sites of origin for Para-testicular RMS: spermatic cord; epididymis; testis and tunica vaginalis. It is noted in 7 % of all patients in IRS and accounts for 17% of all malignant intrascrotal tumours in children less than fifteen years of age [1]. Alveolar; Embryonal; Botryoid embryonal; Spindle cell embryonal; Anaplastic These are according to the internation RMS classification. rRMS is the most frequent variant, encompassing 60% of cases; it is also known as embryonal RMS (eRMS). Para-testicular rhabdomyosarconi is often associated with rather vague clinical symptoms. The diagnostic procedures are – ultrasonography (US) which is used to identify the tumor localization if it cannot be determined by physical examination [2]. For histological diagnosis and the initial treatment, the initial surgical procedure is radical orchiectomy through the inguinal route ligated initially irrespective of the disease state. Like most adult soft tissue sarcomas, rhabdomyosarcoma is chemosensitive; this is why chemotherapy forms a section of the treatment regimen; typical drugs are Actinomycin D, vincristine and cyclophosphamide. Radiotherapy is also sometimes incorporated in this treatment [3].

How to Cite : Hamayun, S., Hidayat, W., Rafi, H., & Nawaz, H. (2024). Recurrent Para-Testicular Rhabdomyosarcoma: A Case Report. Pakistan Journal of Urology (PJU), 2(01), 114–117. <u>https://doi.org/10.69885/pju.v2i01.65</u>

Corresponding Author: Wagma Hidayat

Department of General Surgery, Hayatabad medical complex Peshawar

Email: dr.wagmahidayat707@gmail.com

https://orcid.org/0009-0008-6760-9570

Cell no: +92-313-9636549

1				
(Article History			
	Received:	January	20-2024	
	Revision:	February	16-2024	
	Accepted:	April	22-2024	
	Published:	July	05-2024	

CASE PRESENTATION:

16-year-old boy presented in Surgical OPD with swelling in the right inguinal region of one months duration, who had right radical orchiectomy done one year back for scrotal mass biopsy revealed embryonal rhabdomyosarcoma of para testicular origin while testes and speramtic cord were negative for tumor. He had completed adjuvant chemo-therapy of six cycles of VCR 2 gm, CTX 1 gm and DOX 90 mg. There we saw the scar at the operation site in the right inguinal area and a hard irregular rubbery immobile mass which extended up to the scrotum. Ophthaloscopic examination was also unproductive; transillumination test was also negative. Routine investigations were normal and so was the liver function whereas, tumor markers including LDH, alpha-fetoprotein and beta-HCG were well within normal range. On the abdominal ultrasound there was a lobulated mass of size 5x9 cm in the right inguinal region that had micro-calcification. On the abdomen and pelvic CT scan, a deforming mass of size 10cm X 6 cm was observed in the right inguinal region with fat stranding and enlarged nodes were also noted and multiple indeterminate pulmonary nodules were also observed (Fig 1 & 2). A tru-cut biopsy was performed and immune staining with desmin and myo-D was positive as shown in figures 3 and 4 and the diagnosis of recurrent embryonal rhabdomyosarcoma was made and confirmation was sought from other pathologists. A preoperative patient was on the stage. In course of surgery, a massive necrotic area was seen involving the adductor muscles and the lymph nodes above and below the oblique abdominal muscle; the scrotum was also found to have necrosis. The mass was felt to be misconceived as non-resectable and consequently debulking was done (Figure 5). The specimen was finally subjected to histopathology as the one depicted in the figure 6. He was subsequently referred to a medical oncologist for assessment and management of the case.



Figure 1: Ct Abdomen Pelvis Shows A Large Right Inguinal Region Mass As Pointed By The Red Arrow



Figure 2: Ct Chest Shows Some Indeterminate Pulmonary Infiltrates As Pointed By The Red Arrows



Figure 03: Indeterminate Pulmonary Infiltrates As Pointed By The Red Arrows



Figure 4: Desmin Positive and Myogenin D Positive



Figure 5: Debulked Tissues Of The Tumour

DISCUSSION

Alveolar rhabdomyosarcoma (RMS) serves from embryonal mesencies and is one of the most malignant tumors^[4]. It has four subtypes: These areembryonal, alveolar, pleomorphic and spindle cell. RMS typically arises from soft tissues and affects mainly the head and neck, ocular and orbital region, GU tract and genitals, and limbs[5]. Testicular RMS is also known as para - testicular RMS and is the seventh most prevalent RMS responsible for 7% of all RMS; it is a rather rare tumor that develops from spermatocytic cells of epididymal and spermatic cord origin[6]. This tumor is seen in children and young adults and therefore the age distribution is bimodal with age of 5 and 16 respectively as in the above case. Depending on the stage of development

and it is the most common: the disease can be cured if identified in its early stage. Nevertheless, the relapse rate is quite high, and in our patient, recurrent bilateral pulmonary nodules were diagnosed one year after right orchiectomy. Most para-testicular tumours are found incidentally and are painless mass in the scrotum at the time of they presentation. have often spread hematogenously to iliac and paraaortic lymph nodes or lungs and bones [7]. The oedema is asymptomatic and so patients do not seek medical help and end up presenting at the advanced stage of the disease. The manoeuvres that should be included are the scrotal palpation, inguinal lymph node, other lymph node, general physical examination and the systemic examination for metastasin[8]. Para_testicular RMS is amongst the causes of scrotal swellings and treatment with inguino scrotal ultrasound is useful in differentiating it from other conditionsdue to the demonstration of the lobulated heterogeneous mass that characterizes this tumor vulgaris. Tumor marker assays that are normally elevated in testicular swellings are usually normal in paratesticular RMS although there may be a slight rise in the LDH [9]. For staging of disease and to check loco-regional invasion or metastasis, Thoraco-Abdomino-Pelvic CT and MRI are the gold standard. All these imaging techniques assists in planing the treatment. Treatment of para- testicular RMS should consist of surgery - radical, retroperitoneal lymph node dissection. radiotherapy and chemotherapy[10]. In radical orchiectomy essential histological information is obtained and this is the first operation carried out. This is then followed by chemotherapy, which normally takes 4-6 cycles of a program such as Vincristine, Cyclophosphamide and Adriamycin (VAC), which is known to be effective. Regional

lymph node and metastatic disease is mainly treated by radiotherapy. For the children affected by the paratesticular RMS the prognosis is comparatively better than that of the adults, therefore, early diagnosis and multifaceted treatment is advisable [11]. Such patients therefore require follow up and regular evaluation, in order to note the re occurrences and any consequent effects. It becomes important to rise the level of attention and to detect the disease in its early stages to improve the prognosis and survival rate of the para-testicular R-Med patients.

CONCLUSION

we suspect that the diagnosis of para testicular RMS is not straightforward in many cases and ultrasonography becomes particularly beneficial if physical examination only is inconclusive. The management consists of the surgical treatment that is the removal of the tumor with the affected testicle, orchiectomy; the chemotherapeutic treatment usually Actinomycin D, vincristine, cyclophosphamide and the use of radiotherapy in some special cases.

References:

- 1. Zhu Y, Zhu Z, Xiao Y, Zhu Z. Case report: paratesticular rhabdomyosarcoma. Front Oncol. 2021;11:629878. doi:10.3389/fonc.2021.629878.
- 2. Bouchikhi AA, Mellas S, Tazi MF, et al. Embryonic paratesticular rhabdomyosarcoma: a case report. J Med Case Rep. 2020;14(1):93. doi:10.1186/s13256-020-02409-7.
- **3.** Kühn AL, Scortegagna E, Nowitzki KM, Kim YH. Ultrasonography of the scrotum in adults. Ultrasonography..2020.Jul;39(3):180-197. doi:10.14366/usg.20028.
- Nasrollahi H, Eslahi A, Ariafar A, Ahmed F, Monabati A. Primary rhabdomyosarcoma of kidney with local recurrence and liver metastasis in adults: a case report. J Kidney Cancer VHL. 2022 Apr 14;9(1):55-60. doi:10.15586/jkcvhl.v9i1.192.
- Sadiq AS, Atallah W, Khusid J, Gupta M. The surgical technique of mini percutaneous nephrolithotomy. J Endourol..2021Sep;35(S2):S-68. doi:10.1089/end.2020.0900.
- Kandemir E, Savun M, Sezer A, Erbin A, Akbulut MF, Sarılar Ö. Comparison of miniaturized percutaneous nephrolithotomy and standard percutaneous nephrolithotomy in secondary patients: a randomized prospective study. J Endourol. 2020 Jan;34(1):26-32.

doi:10.1089/end.2019.0573.

- Geraghty RM, Davis NF, Tzelves L, Lombardo R, Yuan C, Thomas K, et al. Best practice in interventional management of urolithiasis: an update from the European Association of Urology Guidelines Panel for Urolithiasis 2022. Eur Urol Focus. 2023 Jan;9(1):199-208. doi:10.1016/j.euf.2022.07.010.
- **8.** Ferrari A, Dileo P, Casanova M, et al. Rhabdomyosarcoma in adults: updated treatment and outcomes from a single institution (2020). Future Oncol. 2021;17(16):2053-2062. doi:10.2217/fon-2020-1149.
- **9.** Amparore D, Campi R, Checcucci E, et al. Forecasting the future of urology practice: review of priority procedures during the COVID-19 pandemic. Eur Urol Focus. 2020 Sep;6(5):1032-1048. doi:10.1016/j.euf.2020.04.014.
- 10. Qin P, Zhang D, Huang T, Fang L, Cheng Y. Comparison of mini percutaneous nephrolithotomy and standard percutaneous nephrolithotomy for renal stones > 2 cm: a systematic review and meta-analysis. Int Braz J Urol. 2022 Jul;48(4):637-648. doi:10.1590/S1677-5538.IBJU.2021.0640.
- **11.** Kallidonis P, Tsaturyan A, Lattarulo M, Liatsikos E. Minimally invasive percutaneous nephrolithotomy (PCNL): techniques and outcomes. Turk J Urol. 2020 Nov;46(Suppl 1):S58-S64. doi:10.5152/tud.2020.20318.

Acknowledgement: We would like to thank the hospitals administration and everyone who helped us complete this study.

Disclaimer: Nil Conflict of Interest: Nil Funding Disclosure: Ni

Authors Contribution:

Concept & Design of Study: Abdul Haseeb1,

Drafting: Sana Hamayun2

Data Analysis: Wagma Hidayat2, Hamza Rafi2

Critical Review: All Mention Authors Approved the Final version



Licensing and Copyright Statement

All articles published in the *Pakistan Journal of Urology* are licensed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0)

This license allows users to **share** (copy and redistribute) and **adapt** (remix, transform, and build upon) the published material for **any purpose, including commercial**, provided appropriate credit is given to the original author(s) and the source (*Pakistan Journal of Urology*), a link to the license is provided, and any changes made are indicated.

This work is licensed under a Creative Commons Attribution 4.0 International License. © The Author(s) 2024